

# Bonvadis<sup>®</sup>

## Topical Cream

### DEVICE DESCRIPTION

Bonvadis Topical Cream is a non-sterile, water based, semi-viscous formulation, for over the counter (OTC) use. Bonvadis Topical Cream contains methyl paraben and propyl paraben to preserve the product before opening and between uses.

The product is indicated for the management of minor skin lacerations, abrasions, cuts, and minor burns. This formulation helps to maintain the wound moist environment. The moist wound environment is conducive for wound healing.

### INGREDIENTS

Purified water, Liquid petrolatum, White petrolatum, Propylene glycol, Cetyl stearyl alcohol, Tween 60, Span 60, *Centella asiatica* extract, Mexican mint fragrance, Methyl paraben, and Propyl paraben.

### INDICATIONS FOR USE

For Over the Counter Use (OTC):

Bonvadis Topical Cream is indicated for the management of minor skin lacerations, abrasions, cuts, and minor burns.

### DIRECTIONS FOR USE

- For topical use only
- Bonvadis Topical Cream is a multiple use wound dressing intended to be applied daily as a primary dressing to a wound.
- Gently cleanse the wound and apply a generous layer of Bonvadis Topical Cream on the surface of the wound.
- Cover Bonvadis Topical Cream with an appropriate secondary dressing, such as gauze.
- The cumulative use of Bonvadis Topical Cream should not exceed 15 g (one tube) in 28 days.

### CONTRAINDICATIONS

Do not use Bonvadis Topical Cream if you have a known sensitivity to any Bonvadis ingredients.  
Do not use for the treatment of third degree burns or to control heavy bleeding.

### WARNINGS AND CAUTIONS


- For external use only.
- Do not use if you have a known sensitivity or allergies to Bonvadis or any of its ingredients.
- Keep out of the reach of children.
- Bonvadis Topical Cream should not be ingested or taken internally. If ingested or taken internally, contact a doctor immediately.
- The product should be discarded after use of 28 days. Safety testing has not been conducted on the product to support the use for more than 28 days.
- Bonvadis Topical Cream does not contain a sunscreen and should not be used prior to extended exposure to the sun.
- Upon observation of the sign of wound infection, consult a healthcare professional and appropriate treatment should be initiated.
- The cumulative use of the subject product is limited to 15 g (one tube) per procedure and used on a single patient only.


### SHELF LIFE AND STORAGE

Bonvadis cannot be used after the expiry date. The expiry date is stated on the package.

Bonvadis Topical Cream should be stored away from sunlight. The storage condition is under ambient temperature and ambient humidity.

### SYMBOL DEFINITION

 Non-sterile

 Use by Date

 Caution, Consult Accompanying Documents

 Batch Number

 Consult Instructions for Use

 Do Not Use if Package is Damaged

 Temperature Limit

 Manufacturer